

Factors of Regular Graphs

MIKIO KANO

Department of Mathematics, Akashi Technological College, Uozumi, Akashi 674, Japan

Communicated by the Managing Editors

Received June 11, 1985

A spanning subgraph F of a graph G is called a $[k-1, k]$ -factor if $k-1 \leq d_F(x) \leq k$ for all vertices x of G , where $d_F(x)$ denotes the degree of x in F . Tutte proved that if r is an odd integer, then every r -regular graph has a $[k-1, k]$ -factor for every integer k , $0 < k < r$. We prove that if r is odd and $0 < k \leq 2r/3$, then every r -regular graph has a $[k-1, k]$ -factor each of whose components is regular. © 1986 Academic Press, Inc.

1. INTRODUCTION

We consider a finite graph G which may have multiple edges but has no loops. A graph without multiple edges is called a *simple graph*. We denote by $V(G)$ and $E(G)$ the set of vertices and the set of edges of G , respectively. We write $d_G(x)$ for the degree of a vertex x in G . Let a , b , and r be integers such that $0 \leq a \leq b$ and $r > 0$. A spanning subgraph F of G is called an $[a, b]$ -factor of G if $a \leq d_F(x) \leq b$ for all $x \in V(G)$, and we usually call an $[r, r]$ -factor an r -factor. An r -regular graph is a graph in which each vertex has degree r . Other notation and definitions not defined in this paper can be found in [3 or 4].

Tutte [12] ([4, p. 77]) proved that for any odd integer r and any integer k ($0 < k < r$), every r -regular graph has a $[k-1, k]$ -factor. It was proved in [6, 11] that every regular graph has a $[1, 2]$ -factor each of whose components is regular. Enomoto and Saito [5] gave the following conjecture: Every r -regular graph has a $[k-1, k]$ -factor each of whose components is regular for any k , $0 < k < r$. Note that this conjecture is true when r is even by Petersen's 2-factorable theorem (see Lemma 1). So the essential part of this conjecture is the case that r is odd. Main results of this paper are the following two theorems, and a theorem on $[a, b]$ -factors is given in Section 3.

THEOREM 1. *Let r and k be positive integers. If $k \leq 2(2r+1)/3$, then every $(2r+1)$ -regular graph has a $[k-1, k]$ -factor each of whose components is regular.*

THEOREM 2. *Let k and r be positive integers. If $2r+2-\sqrt{2r+2} < k \leq 2r$, then there exists a simple $(2r+1)$ -regular graph that has no $[k-1, k]$ -factor each of whose components is regular.*

Some results related to our theorems can be found in a survey article [1].

2. PROOFS OF THEOREMS 1 AND 2

Let G be a graph, and g and f be integer-valued functions defined on $V(G)$ such that $g(x) \leq f(x)$ for all $x \in V(G)$. A spanning subgraph F of G is called a (g, f) -factor if $g(x) \leq d_F(x) \leq f(x)$ for all $x \in V(G)$. A (g, f) -factor satisfying $g(x) = f(x)$ for all $x \in V(G)$ is briefly called an f -factor. For a vertex subset X of G , we write $G - X$ for the subgraph of G obtained from G by deleting the vertices in X together with their incident edges. Similarly, for an edge subset Y of G , $G - Y$ denotes the subgraph of G obtained from G by deleting all the edges in Y . For two disjoint subsets S and T of $V(G)$, we denote by $e_G(S, T)$ or $e(S, T)$ the number of edges of G joining S and T .

LEMMA 1 (Petersen [10]; [3] p. 166). *Every $2r$ -regular graph has a $2k$ -factor for every integer k , $0 < k < r$.*

LEMMA 2 (Lovász [9]). *Let G be a graph, and g and f be integer-valued functions defined on $V(G)$ satisfying $g(x) \leq f(x)$ for all $x \in V(G)$. Then G has a (g, f) -factor if and only if*

$$\delta(S, T) = \sum_{t \in T} (d_G(t) - g(t)) + \sum_{s \in S} f(s) - e_G(S, T) - h(S, T) \geq 0$$

for all disjoint subsets S and T of $V(G)$, where $h(S, T)$ denotes the number of components C of $G - (S \cup T)$ such that $g(x) = f(x)$ for all $x \in V(C)$ and

$$\sum_{x \in V(C)} f(x) + e_G(V(C), T) \equiv 1 \pmod{2}.$$

Lemma 2 plays an important role through this paper, and its short proof is given in [13]. The next lemma is obtained by using Tutte's f -factor theorem [11].

LEMMA 3 [7]. *Let G be an n -edge-connected graph ($n \geq 1$), θ be a real number such that $0 \leq \theta \leq 1$, and f be an integer-valued function defined on $V(G)$. If (1), (2), and (3) hold, then G has an f -factor:*

- (1) $\sum_{x \in V(G)} f(x) \equiv 0 \pmod{2}$.
- (2) $\varepsilon = \sum_{x \in V(G)} |f(x) - \theta d_G(x)| < 2$.
- (3) $\{f(x) \mid x \in V(G)\}$ consists of even numbers, and $m(1 - \theta) \geq 1$, where $m \in \{n, n + 1\}$ and $m \equiv 1 \pmod{2}$.

LEMMA 4 [2]. Let G be a 2-edge-connected $(2r + 1)$ -regular graph, and h be a positive integer. If $2h \leq 2(2r + 1)/3$, then G has a $2h$ -factor. If $(2r + 1)/3 \leq 2h + 1 \leq 2r + 1$, then G has a $(2h + 1)$ -factor. In particular, for every integer k , $0 < k \leq 2r + 1$, G has a $[k - 1, k]$ -factor each component of which is regular.

Note that the above lemma can be proved by Lemma 3.

LEMMA 5. Let G be a 2-edge-connected $[2r, 2r + 1]$ -graph having exactly one vertex w of degree $2r$. Then

- (1) if $0 \leq 2k \leq 2(2r + 1)/3$, then G has a $2k$ -factor; and
- (2) if $(2r + 1)/3 \leq 2k + 1 \leq 2r + 1$, then G has a $[2k, 2k + 1]$ -factor F such that $d_F(w) = 2k$ and $d_F(x) = 2k + 1$ for all $x \in V(G) \setminus \{w\}$.

Proof. We first prove (1). Define a function f on $V(G)$ by $f(x) = 2k$ for all $x \in V(G)$, and put $\theta = 2k/(2r + 1)$. Then G , f , and θ satisfy conditions (1), (2) ($\varepsilon = 2k/(2r + 1) < 1$) and (3) of Lemma 3. Hence G has an f -factor, which is a $2k$ -factor.

We next prove (2). Let F be a $2k$ -factor given in (1). Then the complement $G - E(F)$ is a desired factor of (2).

Let $\{a, b, \dots, n\}$ be a set of integers. A graph G is called an $\{a, b, \dots, n\}$ -graph if $d_G(x) \in \{a, b, \dots, n\}$ for all $x \in V(G)$. Analogously, we define an $\{a, b, \dots, n\}$ -factor. We denote an edge joining a vertex x to a vertex y by xy or yx . The proof of Theorem 1 heavily depends on the next lemma.

LEMMA 6. Let G be a connected $(2r + 1)$ -regular graph with at least two bridges, and k be a positive integer. If $(2r + 4)/3 \leq k \leq 2(2r + 1)/3$, then G has a $[k - 1, k]$ -factor each component of which is regular.

Proof. We first prove that if $(2r + 1)/3 \leq 2k + 1 \leq 2(2r + 1)/3$, then G has a $[2k, 2k + 1]$ -factor each of whose components is regular. Since G has at least two bridges, it follows that for any bridge vw of G , $G - vw$ has at most one 2-edge-connected component. Let $v_1 w_1, \dots, v_n w_n$ be the bridges of G such that $G - v_i w_i$ has exactly one 2-edge-connected component C_i for each i , $1 \leq i \leq n$, where we assume $w_i \in V(C_i)$ and $v_i \in V(G) \setminus V(C_i)$. Put

$$H = G - \bigcup_{i=1}^n (V(C_i) \setminus \{w_i\}),$$

which is the subgraph of G induced by the edge subset $E(G) \setminus \{E(C_1) \cup \cdots \cup E(C_n)\}$. It is immediate that H is a connected $\{2r+1, 1\}$ -graph with end vertices w_1, \dots, w_n .

We shall prove that H has a $\{2k+1, 1, 0\}$ -factor F such that $d_F(x) = 2k+1$ if $d_H(x) = 2r+1$, and $d_F(w) \in \{1, 0\}$ for all end vertices w . Define two functions g and f on $V(H)$ by

$$\begin{aligned} g(x) &= 2k+1 & \text{if } d_H(x) &= 2r+1, \\ &= 0 & \text{otherwise} \end{aligned}$$

and

$$\begin{aligned} f(x) &= 2k+1 & \text{if } d_H(x) &= 2r+1, \\ &= 1 & \text{otherwise.} \end{aligned}$$

Since a (g, f) -factor of H is a required $\{2k+1, 1, 0\}$ -factor, it suffices to show that $\delta(S, T) \geq 0$ for all $S, T \subset V(H)$ such that $S \cap T = \emptyset$ by Lemma 2. Suppose $\delta(S, T) < 0$ for some disjoint subsets S and T of $V(H)$. Since H has at least two end vertices, $\delta(\phi, \phi) = -h(\phi, \phi) = 0$. Hence $S \cup T \neq \emptyset$. Choose disjoint subsets S and T of $V(H)$ in such a way that $|S \cup T|$ is minimum subject to $\delta(S, T) < 0$. Denote by D_1, \dots, D_m the components of $H - (S \cup T)$ which satisfy the conditions on $h(S, T)$ in Lemma 2.

We now prove that neither S nor T contains end vertices of H . Assume S contains an end vertex w . Let wz be the edge of H incident with w . It follows that

$$\begin{aligned} \delta(S, T) &= \sum_{t \in T} (d_H(t) - g(t)) + \sum_{s \in S - w} f(s) + 1 - e(S, T) - h(S, T) \\ &= \delta(S - w, T) + 1 - e(w, T) - h(S, T) + h(S - w, T). \end{aligned}$$

If $z \in T$, then $e(w, T) = 1$ and $h(S, T) = h(S - w, T)$. Thus $\delta(S - w, T) = \delta(S, T) < 0$, which contradicts the choice of S and T . If $z \in S$, then $e(w, T) = 0$ and $h(S, T) = h(S - w, T)$, and so $\delta(S - w, T) = \delta(S, T) - 1 < 0$, a contradiction. If $z \in V(D_i)$ for some D_i , then $e(w, T) = 0$ and $h(S - w, T) = h(S, T) - 1$, and so $\delta(S - w, T) = \delta(S, T) < 0$, a contradiction. If z is contained in a component of $G - (S \cup T)$ that is not D_i ($1 \leq i \leq m$), then $e(w, T) = 0$ and $h(S - w, T) = h(S, T)$. Hence $\delta(S - w, T) < 0$, a contradiction. Therefore S does not contain any end vertex of H .

Suppose T contains an end vertex w . Let wz be the edge of H . Then it follows that

$$\delta(S, T) = \delta(S, T - w) + 1 - e(S, w) - h(S, T) + h(S, T - w).$$

If $z \in T$, then $\delta(S, T - w) = \delta(S, T) - 1$, a contradiction. If $z \in S$, then $\delta(S, T - w) = \delta(S, T)$, a contradiction. If $z \in V(D_i)$ for some D_i , then

$\delta(S, T-w) = \delta(S, T)$, a contradiction. If z is contained in a component of $H - (S \cup T)$ that is not D_i ($1 \leq i \leq m$), then $\delta(S, T-w) = \delta(S, T) - 1$, a contradiction. Therefore T does not contain any end vertices of H . Consequently, $S \cup T$ contains no end vertices of H .

By the conditions on D_i (see Lemma 2), we have that $d_H(x) = 2r + 1$ for all $x \in V(D_i)$. Moreover it follows from the conditions on D_i that

$$\sum_{x \in V(D_i)} f(x) + e(V(D_i), T) \equiv |V(D_i)| + e(V(D_i), T) \equiv 1 \pmod{2}. \quad (1)$$

It is trivial that $e(S \cup T, V(D_i)) \geq 1$ as $S \cup T \neq \emptyset$. If $e(S \cup T, V(D_i)) = 1$, then the bridge joining $S \cup T$ to D_i or a certain bridge of D_i must be chosen in $\{v_1 w_1, \dots, v_n w_n\}$, a contradiction. Therefore

$$e(S \cup T, V(D_i)) \geq 2. \quad (2)$$

Let $h(S, T) = m$ and $\theta = (2r - 2k)/(2r + 1)$. Then $0 < \theta < 1$ and we obtain

$$\begin{aligned} \delta(S, T) &= (2r - 2k) |T| + (2k + 1) |S| - e(S, T) - h(S, T) \\ &= \frac{2r - 2k}{2r + 1} \sum_{t \in T} d_H(t) + \frac{2k + 1}{2r + 1} \sum_{s \in S} d_H(s) - e(S, T) - m \\ &\geq \theta \left\{ e(T, S) + \sum_{i=1}^m e(T, V(D_i)) \right\} \\ &\quad + (1 - \theta) \left\{ e(S, T) + \sum_{i=1}^m e(S, V(D_i)) \right\} - e(S, T) - m \\ &= \sum_{i=1}^m \{ \theta e(T, V(D_i)) + (1 - \theta) e(S, V(D_i)) - 1 \}. \end{aligned}$$

Put

$$\Delta(D_i) = \theta e(T, V(D_i)) + (1 - \theta) e(S, V(D_i)) - 1.$$

We shall show that $\Delta(D_i) \geq 0$ for all i , $1 \leq i \leq m$. If it is proved, then $\delta(S, T) \geq 0$, which is a desired contradiction, and we conclude that H has a (g, f) -factor. If $e(T, V(D_i)) \geq 1$ and $e(S, V(D_i)) \geq 1$, then $\Delta(D_i) \geq 0$. Suppose $e(T, V(D_i)) = 0$. By the congruence expression (1), we have $|V(D_i)| \equiv 1 \pmod{2}$. Since

$$\sum_{x \in V(D_i)} d_H(x) = 2 |E(D_i)| + e(S \cup T, V(D_i)) = (2r + 1) |V(D_i)|, \quad (3)$$

we have $e(S, V(D_i)) \equiv |V(D_i)| \equiv 1 \pmod{2}$. By the inequality (2), we can get $e(S, V(D_i)) \geq 3$. Therefore $\Delta(D_i) \geq (1 - \theta) \times 3 - 1 \geq 0$. Next assume

$e(S, V(D_i))=0$. By (3), we have $e(T, V(D_i)) \equiv |V(D_i)| \pmod{2}$. But this congruence expression contradicts (1). Hence this case does not occur.

Consequently, H has a $\{2k+1, 1, 0\}$ -factor F with the property that $d_F(x)=2k+1$ if $d_H(x)=2r+1$, and $d_F(w) \in \{1, 0\}$ for all end vertices w .

We can now construct a $[2k, 2k+1]$ -factor of G each component of which is regular as follows. Let F be a $\{2k+1, 1, 0\}$ -factor of H given above. If $d_F(w_i)=0$, then we take a $2k$ -factor $R(i)$ of C_i by Lemma 5. If $d_F(w_i)=1$, then we take a $[2k, 2k+1]$ -factor $R(i)$ of C_i such that $d_{R(i)}(w_i)=2k$ and $d_{R(i)}(x)=2k+1$ for all $x \in V(C_i) \setminus \{w_i\}$ by Lemma 5. Then the union $F \cup R(1) \cup \cdots \cup R(n)$ is a $[2k, 2k+1]$ -factor of G with regular components.

We next prove that if $(2r+4)/3 \leq 2k \leq 2(2r+1)/3$, then G has a $[2k-1, 2k]$ -factor each of whose components is regular. Define a subgraph H of G as above. Then H has a $\{2k-1, 1, 0\}$ -factor F' such that $d_{F'}(x)=2k-1$ if $d_H(x)=2r+1$, and $d_{F'}(w) \in \{1, 0\}$ for all end vertices w . By using Lemma 5, we can construct a $[2k-1, 2k]$ -factor of G with regular components from F' in the same way. Consequently, the proof of this lemma is complete.

Proof of Theorem 1. We prove the theorem by induction on $2r+1$. Let G be a $(2r+1)$ -regular graph and k be an integer such that $2 \leq k \leq 2(2r+1)/3$. Note that every regular graph has a $[0, 1]$ -factor with regular components since it has a 0-factor. By Lemma 4, we may assume that G is not 2-edge-connected. Suppose G has exactly one bridge vw . Then each component C of $G-vw$ is a 2-edge-connected $[2r, 2r+1]$ -graph possessing one vertex of degree $2r$. Thus C has a k -factor or a $(k-1)$ -factor by Lemma 5. Therefore G has a k -factor or a $(k-1)$ -factor, and the theorem holds. Consequently, we may assume G has at least two bridges.

By Lemma 6, a 3-regular graph with at least two bridges has a $[1, 2]$ -factor with regular components. Hence every 3-regular graph has a $[1, 2]$ -factor with regular components, and so the theorem is true if $2r+1=3$. Similarly, we can show that every 5-regular graph has a $[2, 3]$ -factor F_1 with regular components. Since each 3-regular component of F_1 has a $[1, 2]$ -factor with regular components, F_1 has a $[1, 2]$ -factor with regular components, which is of course a desired $[1, 2]$ -factor of G . Hence the theorem follows for $2r+1=5$. In general, if a $(2r+1)$ -regular graph G has an $[h-1, h]$ -factor F_2 with regular components and if each component of F_2 has a $[k-1, k]$ -factor with regular components, then G has a $[k-1, k]$ -factor with regular components. By this argument, we can verify that if $2r+1 \leq 17$, then the theorem holds. Suppose $2r+1 \geq 19$. If $(2r+4)/3 \leq k \leq 2(2r+1)/3$, then a $(2r+1)$ -regular graph G has a $[k-1, k]$ -factor with regular components by Lemma 6. Hence we may assume $k < (2r+4)/3$. Let h be the greatest integer not exceeding

$2(2r+1)/3$. Then $h \geq 4r/3$, and G has an $[h-1, h]$ -factor F with regular components. Since $2(h-1)/3 \geq 2(4r-3)/9$ and $2r+1 \geq 19$, we have $2(h-1)/3 \geq (2r+4)/3 > k$. Hence each component of F has a $[k-1, k]$ -factor with regular components by Lemma 1 or by the inductive hypothesis. Therefore G has a $[k-1, k]$ -factor with regular components, and we conclude that the proof of Theorem 1 is complete.

Proof of Theorem 2. Let k and r be positive integers such that $2r+2 - \sqrt{2r+2} < 2k \leq 2r$. Let k' be an odd integer that is one of $\{2k-1, 2k+1\}$ and not equal to $2r+1$. Let K_{2r+3} denote the complete graph with vertex set $\{a_1, \dots, a_{2r+3}\}$. We obtain the graph R from K_{2r+3} by deleting edges $a_1 a_2, a_1 a_3, \dots, a_1 a_{2r-2k+5}, a_{2r-2k+6} a_{2r-2k+7}, \dots, a_{2r+2} a_{2r+3}$. It is clear that $d_R(a_1) = 2k-2$ and $d_R(a_i) = 2r+1$ for all i , $2 \leq i \leq 2r+3$. Let $R(1), \dots, R(2r)$ be copies of R , and let b_i be the vertex of R_i whose degree is $2k-2$ for all i . We construct a graph H with vertex set $V(R(1)) \cup \dots \cup V(R(2r)) \cup \{c_1, \dots, c_{2r-2k+2}, v\}$ as follows. Join every b_i to all c_j ($1 \leq j \leq 2r-2k+2$) and v by new edges, and add new edges $c_1 c_2, c_3 c_4, \dots, c_{2r-2k+1} c_{2r-2k+2}$ (see Fig. 1). Then $d_H(v) = 2r$ and $d_H(x) = 2r+1$ for all $x \in V(H) \setminus \{v\}$.

Let H_1, \dots, H_{2r+1} be copies of H , and let v_i be the vertex of H_i whose degree is $2r$ for every i . We now construct a $(2r+1)$ -regular graph G as follows, which has the required property. Set $V(G) = V(H_1) \cup \dots \cup V(H_{2r+1}) \cup \{w\}$, and join each v_i to w by a new edge. We shall prove that G has no $\{2k, k'\}$ -factor each component of which is regular.

Suppose G has a $\{2k, k'\}$ -factor with regular components. Then it is obvious that some H_i has a $\{2k, k'\}$ -factor with regular components. Thus it suffices to show that H cannot have such a factor.

Assume H has a $\{2k, k'\}$ -factor F with regular components. Since the order of H is odd, F has at least one $2k$ -regular component C of odd order. Suppose $V(C) \cap V(R(i)) \neq \emptyset$ for some i , $1 \leq i \leq 2r$. Since $d_H(b_i) - d_{R(i)}(b_i) = 2r - 2k + 3 < 2k$, C contains at least one vertex in $V(R(i)) \setminus \{b_i\}$, and so $|V(C) \cap V(R(i))| \geq 2k+1$. If C does not contain a vertex x in $V(R(i)) \setminus \{b_i\}$, then $d_{F \setminus C}(x) \leq 2r+3 - (2k+1) < 2k+1$, a contradiction. Hence $V(R(i)) \setminus \{b_i\} \subset V(C)$. If $b_i \notin V(C)$, then C is of even order, a con-




FIG. 1. The graph H .

tradiction. Therefore, $V(C) \cap V(R(i)) \neq \emptyset$ implies that $V(C) \supset V(R(i))$ and that b_i is joined to at least one vertex, say u , in $\{c_1, \dots, c_{2r-2k+2}, v\}$ by an edge of C (since $d_{R(i)}(b_i) = 2k - 2 < d_C(b_i)$). For each vertex u_2 in $\{c_1, \dots, c_{2r-2k+2}, v\} \setminus \{u_1\}$, we have $d_F(u_1) + d_F(u_2) \geq 4k - 1 > 2r + 2$, and so u_1 and u_2 are adjacent to a common b_i . Hence $\{c_1, \dots, c_{2r-2k+2}, v\} \subset V(C)$. If $V(C) \cap V(R(j)) = \emptyset$ for some j , $1 \leq j \leq 2r$, then $R(j)$ contains a $2k$ -regular component C' of F since $R(j)$ is of odd order. By the same argument as above, it follows that $V(C') \supset V(R(j))$ and b_j is joined to a vertex in $\{c_1, \dots, c_{2r-2k+2}, v\}$ by an edge of C' , which is contrary to $\{c_1, \dots, c_{2r-2k+2}, v\} \subset V(C)$. Consequently, C is equal to F , in particular, F is a $2k$ -factor of H .

On the other hand, we can show that H has no $2k$ -factor by Lemma 2, which establishes the proof of Theorem 2. Define two functions g and f on $V(H)$ by $g(x) = f(x) = 2k$ for all $x \in V(H)$, and set $S = \emptyset$ and $T = \{c_1, \dots, c_{2r-2k+2}, v\}$. Since $R(1), \dots, R(2r)$ are the components of $H - (S \cup T)$ that satisfy the conditions on $h(S, T)$ in Lemma 2, we obtain

$$\begin{aligned} \delta(S, T) &= (2r + 1 - 2k)(|T| - 1) + 2r - 2k - h(S, T) \\ &= (2r + 1 - 2k)(2r - 2k + 3) - 2r - 1 < 0 \end{aligned}$$

as $2r + 2 - \sqrt{2r + 2} < 2k$. Therefore H has no $2k$ -factors.

3. $[a, b]$ -FACTORS OF GRAPHS

Katerinis [8] recently obtained the following result. Let m, n , and k be positive odd integers such that $m < k < n$. If a graph G has an m -factor and an n -factor, then G has a k -factor. We give a similar result on $[a, b]$ -factors. Note that the inequality (4) in Theorem 3 is equivalent to the condition that the point (m, n) in the plane lies above the straight line passing through the points (a, b) and (c, d) .

THEOREM 3. *Let a, b, c, d, m, n be integers satisfying $0 \leq a < m < c$, $a \leq b, c \leq d$, and $m + 1 \leq n$. Suppose a graph G has both an $[a, b]$ -factor and a $[c, d]$ -factor. If*

$$(n - b)(m - c) \leq (n - d)(m - a), \quad (4)$$

then G has an $[m, n]$ -factor.

Proof. By Lemma 2 with $g(x) = m$ and $f(x) = n$, it suffices to show $\delta(S, T) \geq 0$ for all disjoint subsets S and T of $V(G)$, where

$$\delta(S, T) = \sum_{t \in T} (d_G(t) - m) + n|S| - e(S, T).$$

Note that $h(S, T) = 0$ as $m + 1 \leq n$. Suppose $\delta(S, T) < 0$ for some $S, T \subset V(G)$, $S \cap T = \emptyset$. Since G has an $[a, b]$ -factor, we have

$$\delta_1(S, T) = \sum_{t \in T} (d_G(t) - a) + b |S| - e(S, T) - h_1(S, T) \geq 0.$$

Similarly, since G has a $[c, d]$ -factor, we have

$$\delta_2(S, T) = \sum_{t \in T} (d_G(t) - c) + d |S| - e(S, T) - h_2(S, T) \geq 0.$$

Then we obtain

$$0 > \delta(S, T) - \delta_1(S, T) \geq (a - m) |T| + (n - b) |S| \quad (5)$$

and

$$0 > \delta(S, T) - \delta_2(S, T) \geq (c - m) |T| + (n - d) |S|. \quad (6)$$

If $S = \emptyset$, then $(c - m) |T| < 0$ from (6), a contradiction. Hence $S \neq \emptyset$. By (5) and (6), we get

$$\frac{n - b}{m - a} < \frac{|T|}{|S|} \quad \text{and} \quad \frac{|T|}{|S|} < \frac{n - d}{m - c}$$

and thus $(n - b)(m - c) > (n - d)(m - a)$, which is contrary to (4). Consequently, $\delta(S, T) \geq 0$ for all $S, T \subset V(G)$, $S \cap T = \emptyset$. Thus G has an $[m, n]$ -factor.

COROLLARY. *Let $0 < k < r$. Then every $[r - 1, r]$ -graph has a $[k - 1, k]$ -factor.*

Proof. Put $a = b = 0$, $c = r - 1$, $d = r$, $m = k - 1$, and $n = k$. Then (4) in Theorem 3 follows. Hence the corollary holds.

ACKNOWLEDGMENT

The author wishes to express his thanks to Professor H. Enomoto for his helpful suggestions.

REFERENCES

1. J. AKIYAMA AND M. KANO, Factors and factorizations of graphs—A survey, *J. Graph Theory* **9** (1985), 1–42.
2. F. BÄBLER, Über die Zerlegung regulären Streckenkomplexe ungerader Ordnung, *Comment Math. Helv.* **10** (1938), 275–287.

3. M. BEHZAD, G. CHARTRAND, AND L. LESNIAK-FOSTER, "Graphs & Digraphs," Prindle, Weber & Schmidt, Boston, 1979.
4. B. BOLLABÁS, "Extremal Graph Theory," Academic Press, London, 1978.
5. H. ENOMOTO AND A. SAITO, private communication.
6. A. HAJNAL, A theorem on k -saturated graphs, *Canad. J. Math.* **17** (1965), 720–772.
7. M. KANO, $[a, b]$ -factorization of a graph, *J. Graph Theory* **9** (1985), 129–146.
8. P. KATERINIS, Some conditions for the existence of f -factors, *J. Graph Theory* **9** (1985), 513–521.
9. L. LOVÁSZ, Subgraphs with prescribed valencies, *J. Combin. Theory* **8** (1970), 391–416.
10. J. PETERSEN, Die Theorie der regulären Graph, *Acta Math.* **15** (1891), 193–220.
11. W. T. TUTTE, The factors of graphs, *Canad. J. Math.* **4** (1952), 314–328.
12. W. T. TUTTE, The subgraph problem, *Ann. Discrete Math.* **3** (1978), 289–295.
13. W. T. TUTTE, Graph factors, *Combinatorica* **1** (1981), 79–97.